

HOW TO ADMINISTER SOLU-CORTEF®

(hydrocortisone sodium succinate for injection, USP)

via INTRAMUSCULAR INJECTION using the ACTO-VIAL® System

Introduction



- Education and training are paramount for the proper technique of intramuscular injections
 - This training may be used to prepare a person for self-injection, or to train a caregiver, spouse or other person who may need to inject Solu-Cortef® using an Acto-vial® during an emergency
- In addition to preparation before an injection is needed, staying calm and focused will help you to successfully administer Solu-Cortef® to yourself or your loved one

Reminder: some symptoms of an adrenal crisis may vary slightly by individual, be sure to discuss with your doctor what symptoms you should look for to indicate a risk of a crisis and when you should inject Solu-Cortef®

General Preparation



- Wash your hands with soap and water or an alcohol hand sanitizer
- Gather your equipment*
 - Needles (2)
 - Syringe
 - Alcohol pad
 - Bandaid



^{*}NADF recommends that you keep all of these items in an emergency kit along with the 1-page instruction handout version of this presentation

Preparation of ACTO-VIAL®



- 1. Double check expiration date on side of the vial*
- 2. Push the stopper with your thumb to engage the powdered medication with the solution in the vial
- 3. Mix the power and solution
 - *DO NOT SHAKE VIAL*
 - Gently "swirl" the vial or rock the vial back and forth to allow it to mix completely






^{*}Please follow FDA recommendations regarding extended expiration dates for certain lots for Solu-Cortef ®

Prepare the Injection



- 1. Confirm the medication is completely mixed
- 2. Remove yellow "tab" at the top of the vial
- 3. Wipe top of the vial with an alcohol prep pad





Prepare Syringe



1. Open the package and take out the syringe with needle

■ Your doctor should provide you with guidance about the appropriate needle size and prescribe a needle typically in the range of 22 - 25 gauge, 1 to 1.5 inches in length

2. Draw air into syringe

Insert needle/syringe completely through the rubber stopper. Invert the vial and and gently insert the air from the syringe





Withdraw the medication



- Invert the bottle and gently withdraw the correct dose*
- After removal, place the cap gently back on the syringe and remove any air bubbles by "tapping" the side of the syringe and expel any excess air
- Preferably replace the needle on the syringe with a new needle




^{*2} ml for adults, a lesser amount may be required for children. Please check with your Pediatric Endocrinologist for dosing guidance

Identify the Injection Site



- It is recommended you identify this site in partnership with your doctor and based upon your unique body shape/size prior to needing an emergency injection
 - Since most people will be able to use the thigh for an injection, NADF has detailed those instructions here
- Visualize your thigh in thirds
 - Use your hands to mark the upper third close to your hips and the lower third just above the knee; marker used in photos for illustrative purposes only
- Identify an injection site in the middle "third" on the outer portion of your thigh



Prepare to Inject

National Adrenal Diseases Foundation

- 1. Cleanse the site with an alcohol pad prior to injecting
- 2. Hold the site taut, and holding the syringe similar to a dart, push the needle into the muscle at a 90° angle
 - The injection can be given straight down, or from the side (as pictured)
 - The needle should be pushed in at least ¾ of its length





Administer Injection



- Briskly push the plunger in to administer the medication into the muscle.
- Once injection is complete, withdraw the syringe/needle quickly and apply pressure to the injection site and apply a Band-Aid to the site, if needed

Additional Information



- It is common to feel anxious or uneasy when giving or receiving an injection
- Education, training and preparation is key for self administration of an intramuscular injection
- Remember to call your provider or go to the nearest Emergency Room after you have had an injection of Solu-Cortef®